

12.2. EVIDENCE BASE:

Should AUXILIARY NURSES initiate and maintain injectable contraceptives using a standard syringe?

Problem: Poor access to contraception

Option: Auxiliary nurses initiating and maintaining injectable

contraceptives using a standard syringe

Comparison: Care delivered by other cadres or no care

Setting: Community/primary health care settings in LMICs with poor

access to health professionals

CRITERIA JUDGEMENT EVIDENCE	COMMENTS AND QUERIES
Are the anticipated desirable effects large? No Probably Uncertain Probably yes Varies yes A systematic review (Polus 2012a) searched for studies that assessed the effects and safety of ta	task shifting
Are the anticipated undesirable effects small? Are the anticipated undesirable effects small? No Probably no probably yes per probably yes p	ne delivery of the effects of efore
What is the certainty of the anticipated effects? What is the certainty of the Are the	ganda in evices). It is
Are the desirable effects large relative to the undesirable effects? Are the desirable effects? Safety and patient satisfaction because the quality of the evidence from this study is very low. Annex: page 15 (Oladapo 2012)	
Main resource requirements	
Resource Settings in which auxiliary nurses already provide other care	
Are the resources required small? Are the resources required small? No Probably no Uncertain Probably yes Varies Jes Uncertain Probably yes Varies Incontraceptive methods and promotion; universal precautions Supervision and Regular supervision by midwife or nurse	
Supervision and Regular supervision by midwife or nurse monitoring	
Supplies Injectable contraceptives, syringes, sterile solution, robust supply chain	



	CRITERIA	JUDGEMENT	EVIDENCE	COMMENTS AND QUERIES
	Is the incremental cost small relative to the benefits?	No Probably Uncertain Probably Yes Varies no yes	Uncertain as there is insufficient evidence on effectiveness	
ACCEPTABILITY	Is the option acceptable to most stakeholders?	No Probably Uncertain Probably Yes Varies no yes	We are not aware of any systematic reviews that considered the acceptability of auxiliary nurse interventions. We are therefore uncertain about the acceptability of this intervention to key stakeholders. Indirect evidence: A systematic review (Rashidian 2012) exploring factors that influence the success of doctor-nurse substitution suggests that the acceptability of this intervention to key stakeholders may be mixed: Nurses may be motivated to take on new tasks by increased recognition and job satisfaction (moderate certainty evidence) Doctors may welcome the contribution of nurses where it reduces doctors' workloads (moderate certainty evidence). They may welcome the transfer of certain repetitive tasks to nurses (e.g. pap smears) and nurses seem to be happy with these tasks However, a lack of clarity about nurse roles and responsibilities in relation to other health workers may be a challenge (low certainty evidence) A review of country case studies of task shifting for family planning (Polus 2012b), which mainly included LHW programmes, suggests that recipients appreciate the easy access that community-based or home-based provision of contraceptives provides and appreciate the use of female health workers in the delivery of contraceptives. However, the review also suggests that some health workers may introduce their own criteria when determining who should receive contraceptives, including criteria tied to the recipient's marital status and age. Other factors that may affect the uptake of the intervention are primarily tied to the contraceptives themselves rather than the use of specific types of health workers, including a lack of knowledge about different methods of contraception; religious and other beliefs regarding family planning; a fear of side effects, service fees; and a lack of support from husbands. Annex: page 43 (Rashidian 2012); page 63 (Polus 2012b)	
FEASIBILITY	Is the option feasible to implement?	No Probably Uncertain Probably Yes Varies yes	The intervention requires very few supplies (injectable contraceptives, syringes, sterile solution). However, changes to drug supplies may be needed and the intervention is also likely to require changes to norms or regulations. Training, including in communication about family planning, and supervision is necessary. However, a review of country case studies of task shifting for family planning (Polus 2012b) suggests that auxiliary nurses lacked confidence in their skills, partly because they had insufficient opportunities to practice these skills in settings were demand was low. In addition, systematic reviews of lay health worker, nurse and midwife programmes suggest that sufficient training and supervision is often lacking (moderate certainty evidence) (Glenton, Colvin 2012, Rashidian 2012, Colvin 2012). Annex: page 63 (Polus 2012b); page 26 (Glenton, Colvin 2012); page 20 (Colvin 2012); page 43 (Rashidian 2012)	