

12.2. RECOMMENDATIONS:

Should AUXILIARY NURSE MIDWIVES initiate and maintain injectable contraceptives using a standard syringe?

Problem: Poor access to contraception

Option: Auxiliary nurse midwives initiating and maintaining injectable

contraceptives using a standard syringe

Comparison: Care delivered by other cadres or no care

Setting: Community/primary health care settings in LMICs with poor access to

health professionals

Recommendation	We recommend against the option	We suggest considering the option with targeted monitoring and evaluation	We recommend the option
			\square
	We recommend the use of auxiliary nurse midwives to deliver injectable contraceptives using a standard syringe.		
Justification	There is insufficient evidence on the effectiveness of this intervention. However, this intervention may be a cost-effective, acceptable and feasible approach to making injectable contraceptives available more widely. In addition, the delivery of injections is part of auxiliary nurse midwife practice in a number of settings.		
Implementation considerations	The following should be considered when using auxiliary nurse midwives to deliver injectable contraceptives: The relevant professional bodies should be involved in the planning and implementation of the intervention to ensure acceptability among affected health workers The distribution of roles and responsibilities between auxiliary nurse midwives and other health workers needs to be made clear, including through regulations and job descriptions Changes in regulations may be necessary to support any changes in auxiliary nurse midwives' scope of practice Implementation needs to be in the context of a comprehensive remuneration scheme, in which salaries or incentives reflect any changes in scope of practice. Giving incentives for certain tasks but not for others may negatively affect the work that is carried out Supplies need to be secure Responsibility for supervision needs to be clear and supervision needs to be regular and supportive Because of the sensitivity of sexual and contraceptive issues, planners should consider whether health workers promoting or delivering reproductive health services to women should also be women. It may also be an advantage to ensure that relevant training of female health workers is carried out by females Auxiliary nurse midwives and their supervisors need to receive appropriate initial and ongoing training, including in communicating with recipients and in side effects of different contraceptive methods. Training needs to reinforce that auxiliary nurses should avoid introducing their own criteria for determining who should receive contraception are confidential.		
Monitoring and evaluation	Implementation should include monitoring of the	standard of counseling on contraceptive choices.	
Research priorities			