

The clinical effectiveness and cost-effectiveness of treatments for children with idiopathic steroid-resistant nephrotic syndrome: a systematic review

JL Colquitt,^{1*} J Kirby,¹ C Green,¹ K Cooper¹
and RS Trompeter²

¹ Southampton Health Technology Assessments Centre,
University of Southampton, UK

² Great Ormond Street Hospital NHS Trust, London, UK

* Corresponding author



Executive summary

Health Technology Assessment 2007; Vol. 11: No. 21

Health Technology Assessment
NHS R&D HTA Programme
www.hta.ac.uk





INAHTA

How to obtain copies of this and other HTA Programme reports.

An electronic version of this publication, in Adobe Acrobat format, is available for downloading free of charge for personal use from the HTA website (<http://www.hta.ac.uk>). A fully searchable CD-ROM is also available (see below).

Printed copies of HTA monographs cost £20 each (post and packing free in the UK) to both public **and** private sector purchasers from our Despatch Agents.

Non-UK purchasers will have to pay a small fee for post and packing. For European countries the cost is £2 per monograph and for the rest of the world £3 per monograph.

You can order HTA monographs from our Despatch Agents:

- fax (with **credit card** or **official purchase order**)
- post (with **credit card** or **official purchase order** or **cheque**)
- phone during office hours (**credit card** only).

Additionally the HTA website allows you **either** to pay securely by credit card **or** to print out your order and then post or fax it.

Contact details are as follows:

HTA Despatch
c/o Direct Mail Works Ltd
4 Oakwood Business Centre
Downley, HAVANT PO9 2NP, UK

Email: orders@hta.ac.uk
Tel: 02392 492 000
Fax: 02392 478 555
Fax from outside the UK: +44 2392 478 555

NHS libraries can subscribe free of charge. Public libraries can subscribe at a very reduced cost of £100 for each volume (normally comprising 30–40 titles). The commercial subscription rate is £300 per volume. Please see our website for details. Subscriptions can only be purchased for the current or forthcoming volume.

Payment methods

Paying by cheque

If you pay by cheque, the cheque must be in **pounds sterling**, made payable to *Direct Mail Works Ltd* and drawn on a bank with a UK address.

Paying by credit card

The following cards are accepted by phone, fax, post or via the website ordering pages: Delta, Eurocard, Mastercard, Solo, Switch and Visa. We advise against sending credit card details in a plain email.

Paying by official purchase order

You can post or fax these, but they must be from public bodies (i.e. NHS or universities) within the UK. We cannot at present accept purchase orders from commercial companies or from outside the UK.

How do I get a copy of HTA on CD?

Please use the form on the HTA website (www.hta.ac.uk/htacd.htm). Or contact Direct Mail Works (see contact details above) by email, post, fax or phone. *HTA on CD* is currently free of charge worldwide.

The website also provides information about the HTA Programme and lists the membership of the various committees.



Executive summary

Background

Nephrotic syndrome is a collection of signs and symptoms, including protein in the urine, low blood protein levels, high cholesterol levels and swelling. First line treatment is with oral corticosteroids, but some children do not respond to this treatment. The optimal treatment of steroid-resistant nephrotic syndrome (SRNS) is uncertain.

Objectives

The objectives of this review were to assess the clinical effectiveness and cost-effectiveness of treatments for children with idiopathic SRNS.

Methods

Data sources

Electronic databases were searched from inception to February 2006. Bibliographies of included studies and related papers were checked for relevant studies. Experts were contacted for advice and peer review and to identify additional studies.

Study selection

Titles and abstracts were screened for eligibility by one reviewer and checked by a second. Inclusion criteria were applied to the full text of selected papers by two reviewers, with differences resolved through discussion. Inclusion criteria were:

- intervention: high-dose steroids, immunosuppressive agents, alkylating agents, plasma exchange therapy, angiotensin-converting enzyme inhibitors or fish oils
- patients: children aged 1–18 years with idiopathic SRNS
- studies: systematic reviews of randomised controlled trials (RCTs), RCTs, controlled clinical trials, prospective cohort studies with concurrent controls and economic evaluations; abstracts were considered if sufficient information was presented; non-English-language studies were excluded
- outcomes: remission rates, relapse rates, renal function, adverse effects, long-term renal survival, quality of life, costs and cost-effectiveness.

Data extraction and quality assessment

Data extraction and quality assessment were undertaken by one reviewer and checked by a second, with differences resolved through discussion. The quality of included studies was assessed using criteria from the NHS Centre for Reviews and Dissemination.

Data synthesis

The clinical effectiveness data were synthesised through a narrative review with full tabulation of results. Meta-analysis was undertaken, where appropriate, using the random effects model. Where data allowed, subgroup analysis was undertaken according to renal histopathology [e.g. minimal change nephrotic syndrome (MCNS), focal segmental glomerulosclerosis (FSGS)].

Results

Number and quality of studies

Two systematic reviews and 11 trials were included in the systematic review of clinical effectiveness. The quality of reporting and methodology of the included studies was generally poor. No economic evaluations were identified.

Summary of benefits and risks

No statistically significant difference in remission rates was found between cyclophosphamide plus prednisone and prednisone alone for all children [relative risk (RR) 1.15, 95% confidence interval (CI) 0.65 to 2.05] or those with FSGS (RR 1.01, 95% CI 0.43 to 2.37). Time to response was statistically significantly less with cyclophosphamide [38.4 days (range 6–80) versus 95.5 days (range 61–129), $p < 0.05$]. Death occurred in five patients. Remission rates were not statistically significantly different between intravenous and oral cyclophosphamide. Vomiting was common with intravenous cyclophosphamide, while pneumonia and alopecia occurred in the oral group.

Ciclosporin statistically significantly increased the number of children (both MCNS and FSGS included) with complete remission compared with placebo or supportive treatment (RR 7.66, 95% CI 1.06 to 55.34), but not for the FSGS subgroup (RR 5.83, 95% CI 0.75 to 45.09). One trial

did not contribute to the summary statistic as no patient in either group had remission. One study reported no major side-effects. Adverse effects including infection and hypertension differed little between groups.

No differences were found between azathioprine and placebo, with about 13% of each group having remission.

Complete or partial remission occurred in six out of seven patients on the 18-month methylprednisolone regimen and three out of five patients on the 6-month regimen. Renal function improved in both groups. Adverse events such as hypertension and frequent infections occurred in both groups. One death occurred.

Intravenous dexamethasone and methylprednisolone produced similar complete remission rates (35.1%, 95% CI 22.9 to 48.9, versus 33.3%, 95% CI 14.6 to 46.9) and partial remission rates (12.3%, 95% CI 5.0 to 23.7 versus 14.3%, 95% CI 3.0 to 36.3). Median time to response (about 10 days) and total number of adverse events were also similar. The most common adverse event was hypertension. There was a statistically significant decrease in median urine to albumin creatinine ratio in both groups.

Six-hour urinary albumin and urinary albumin to creatinine ratio decreased statistically significantly with high-dose but not low-dose enalapril. The difference in the urine albumin to creatinine ratio reduction percentage between the two groups was statistically significant in the period before cross-over only. A small number of patients experienced dry cough.

Tuna fish oil was not associated with any statistically significant improvements in proteinuria, creatinine clearance, serum creatinine or lipid profiles compared with placebo.

Summary of costs

A very limited literature was found on costs associated with SRNS in children. Costs consisted of treatment costs, longer term monitoring and management costs, and longer term costs for patients who progress to end-stage renal failure. The pharmaceutical cost of treatment varied considerably: an 8-week course of cyclophosphamide cost less than £6, while a course of ciclosporin cost almost £900 per year. Treatment with tacrolimus, an alternative to

ciclosporin, was estimated to cost in excess of £3400 per year. In addition to pharmaceutical costs, healthcare medical management costs were estimated; varying by treatment strategy, they ranged from £250 to £930 per year in patients not experiencing complications. Other longer term costs may be incurred; these may comprise the cost of care for longer term side-effects and complications, and costs associated with the onset and management of renal failure. Children who fail to respond to treatment are at high risk of developing end-stage renal failure, the costs of which are considerable.

Summary of cost-effectiveness

No published evidence on the cost-effectiveness of treatments for SRNS in children was identified. Subsequent searches were undertaken to identify economic evaluations and economic evidence for SRNS in adults. The current data are sparse and the modelling of the cost-effectiveness of current treatments for SRNS is not feasible at present. It is clear that in future cost-effectiveness analysis it would be inappropriate to compare interventions with a 'no treatment' alternative. It is suggested here that ciclosporin be used as the comparator strategy in future cost-effectiveness analysis, and that the appropriate patient group for analysis may be those patients either not indicated for cyclophosphamide treatment and/or not responding to cyclophosphamide, who would typically be treated with ciclosporin. Where appropriate data on clinical effectiveness were available, a framework for the assessment of the cost-effectiveness of treatment for SRNS was identified in the current review.

Conclusions

Implications for healthcare

The clinical effectiveness literature on treatments for idiopathic SRNS in children is very limited. The available evidence suggests a beneficial effect of ciclosporin on remission rates and of cyclophosphamide on time to remission; however, the strength of the conclusions drawn is limited by the poor quality of the included studies. The other treatments included in this review were each evaluated by only one study, and none found a statistically significant effect. There is insufficient evidence to determine whether or not there is a clinically significant difference. No economic evaluations were identified. The available data on costs and outcomes are sparse and do not permit the

reliable modelling of the cost-effectiveness of treatments for SRNS at present. A modelling framework is suggested, should more relevant data become available.

Recommendation for future research

A well-designed adequately powered RCT comparing ciclosporin with other treatments in children with SRNS without genetic mutation is required.

Publication

Colquitt JL, Kirby J, Green C, Cooper K, Trompeter RS. The clinical effectiveness and cost-effectiveness of treatments for children with idiopathic steroid-resistant nephrotic syndrome: a systematic review. *Health Technol Assess* 2007;**11**(21).

NIHR Health Technology Assessment Programme

The Health Technology Assessment (HTA) programme, now part of the National Institute for Health Research (NIHR), was set up in 1993. It produces high-quality research information on the costs, effectiveness and broader impact of health technologies for those who use, manage and provide care in the NHS. 'Health technologies' are broadly defined to include all interventions used to promote health, prevent and treat disease, and improve rehabilitation and long-term care, rather than settings of care.

The research findings from the HTA Programme directly influence decision-making bodies such as the National Institute for Health and Clinical Excellence (NICE) and the National Screening Committee (NSC). HTA findings also help to improve the quality of clinical practice in the NHS indirectly in that they form a key component of the 'National Knowledge Service'.

The HTA Programme is needs-led in that it fills gaps in the evidence needed by the NHS. There are three routes to the start of projects.

First is the commissioned route. Suggestions for research are actively sought from people working in the NHS, the public and consumer groups and professional bodies such as royal colleges and NHS trusts. These suggestions are carefully prioritised by panels of independent experts (including NHS service users). The HTA Programme then commissions the research by competitive tender.

Secondly, the HTA Programme provides grants for clinical trials for researchers who identify research questions. These are assessed for importance to patients and the NHS, and scientific rigour.

Thirdly, through its Technology Assessment Report (TAR) call-off contract, the HTA Programme commissions bespoke reports, principally for NICE, but also for other policy-makers. TARs bring together evidence on the value of specific technologies.

Some HTA research projects, including TARs, may take only months, others need several years. They can cost from as little as £40,000 to over £1 million, and may involve synthesising existing evidence, undertaking a trial, or other research collecting new data to answer a research problem.

The final reports from HTA projects are peer-reviewed by a number of independent expert referees before publication in the widely read monograph series *Health Technology Assessment*.

Criteria for inclusion in the HTA monograph series

Reports are published in the HTA monograph series if (1) they have resulted from work for the HTA Programme, and (2) they are of a sufficiently high scientific quality as assessed by the referees and editors.

Reviews in *Health Technology Assessment* are termed 'systematic' when the account of the search, appraisal and synthesis methods (to minimise biases and random errors) would, in theory, permit the replication of the review by others.

The research reported in this monograph was commissioned by the HTA Programme as project number 05/37/01. The contractual start date was in September 2005. The draft report began editorial review in March 2006 and was accepted for publication in January 2007. As the funder, by devising a commissioning brief, the HTA Programme specified the research question and study design. The authors have been wholly responsible for all data collection, analysis and interpretation, and for writing up their work. The HTA editors and publisher have tried to ensure the accuracy of the authors' report and would like to thank the referees for their constructive comments on the draft document. However, they do not accept liability for damages or losses arising from material published in this report.

The views expressed in this publication are those of the authors and not necessarily those of the HTA Programme or the Department of Health.

Editor-in-Chief:

Professor Tom Walley

Series Editors:

Dr Aileen Clarke, Dr Peter Davidson, Dr Chris Hyde,
Dr John Powell, Dr Rob Riemsma and Dr Ken Stein

Managing Editors:

Sally Bailey and Sarah Llewellyn Lloyd

ISSN 1366-5278

© Queen's Printer and Controller of HMSO 2007

This monograph may be freely reproduced for the purposes of private research and study and may be included in professional journals provided that suitable acknowledgement is made and the reproduction is not associated with any form of advertising.

Applications for commercial reproduction should be addressed to: NCCHTA, Mailpoint 728, Boldrewood, University of Southampton, Southampton SO16 7PX, UK.

Published by Gray Publishing, Tunbridge Wells, Kent, on behalf of NCCHTA.

Printed on acid-free paper in the UK by St Edmundsbury Press Ltd, Bury St Edmunds, Suffolk.